Types of evolution

- Divergent
- Convergent
- Parallel
- *Co-evolution

The main mechanisms of evolution Also known as the "forces of evolution"

1 . Natural selection

Traits that provide a *reproductive advantage* tend to increase in frequency in a given population over time, while traits that leave individuals at a reproductive disadvantage tend to decrease

Natural selection probably accounts for more changes in allele frequency than any other microevolutionary process

• It occurs when environmental conditions affect the ability of individuals to survive or reproduce depending on their genotype



1. Natural selection

•All populations have the reproductive capacity to increase in size, thus threatening their own survival, forcing a competition for limited resources.

•Individuals of a population share in the same gene pool, but differ in phenotypic details.

•Some alleles promote survival and reproduction and therefore increase in frequency over generations resulting in individuals that differ in one or more heritable traits (evolution).

•There is a difference in the survival and reproductive success of different phenotypes.

Three possible outcomes for natural selection:


- A shift in the range of values for a given trait in some direction.
- Stabilization of an existing range of values.
- Disruption of an existing range of values.

. Natural selection

 $\boldsymbol{\cdot}$ It acts directly on phenotypes and indirectly on genotypes

• Over time, the alleles that produce the most successful phenotypes will increase in the population.

- Less successful alleles will become less common.
- $\boldsymbol{\cdot}$ Change leads to increased fitness and increased adaptation to environment.




- Mutations need not change gene expression, or have an effect on phenotype; because of the redundancy of the genetic code
- Mutations can alter both the DNA and the protein it expresses
 - <u>Neutral</u> have no effect on the viability or reproduction of the organism
 - <u>Beneficial</u> meaning that they are advantageous to the organism
 - $\frac{harmful \ or \ lethal}{protein \ is \ detrimental \ to \ the \ organism}$ if the change they cause in the

The main mechanisms of evolution...

3. Genetic drift

The random fluctuation in allele frequencies over time, due to chance occurrences alone. It is more significant in small populations.

the increase of the other allele is not because it is better adapted (or "fittest") but simply random chance



3. Genetic drift

Sampling error explains why the chance of any given allele becoming more or less prevalent is more pronounced in a small population.

Fixation means that one kind of allele remains at a specified locus in a population.

There are two forms of genetic drift

The founder effect:

A small number of individuals break off and form a separate population . The ratio of alleles in the new population may be very different from the larger previous population.

Genetic bottlenecks:

When populations (or species) are reduced to a very small number such that the genetic diversity of the small population may be much less than the previous population

Often endangered over fished species are reduced to very small numbers

Inbreeding


Inbreeding refers to the nonrandom mating among closely related individuals, which have many alleles in common.

Inbreeding is a form of genetic drift.

It leads to more homozygosity.

It can lower fitness when deleterious recessive alleles are expressed.


Bottlenecks and inbreeding are especially deleterious to endangered species (**restocking**).





Gene flow

is the physical flow of alleles into a population. It tends to keep the gene pools of populations similar. It counters the differences that result from mutation, natural selection, and genetic drift.

Genes move with the individuals when they move out of, or into, a population. The physical flow (and resultant shuffling) tends to minimize genetic variation between populations.







Summary

- 1. All life forms (species) have developed from other species.
- All living things are related to one another to varying degrees through common decent (share common ancestors).
- 3. All life on Earth has a common origin. On other words, that in the distant past, there once existed an original life form and that this life form gave rise to all subsequent life forms.
- 4. The process by which one species evolves into another involves random heritable genetic mutations (change), some of which are more likely to spread and persist in a gene pool than others.